

APPLICATION FOR PATENT

5 Inventor: Shamay Izhar

10 Title: EXOGENIC ALLELISM

15 This is a continuation-in-part of U.S. Provisional Pat. Application
No. 60/151,627, filed August 31, 1999.

20

FIELD AND BACKGROUND OF THE INVENTION

25 The present invention relates to a novel concept in genetic engineering, according to which a eukaryotic organism is transformed and crossed so as to provide an offspring including a first exogene in a first chromosome of a chromosome pair and a second exogene in a second chromosome of the chromosome pair, the exogenes exhibiting allelic relationship, such that the exogenes obligatorily segregate to different gametes. The present invention further relates to expression cassettes for implementing the novel concept, to methods of

implementing same for the obtainment of reversible male sterility in plants, and further to plants and plant products obtained thereby.

The gene is the basic functional unit of heredity. The gene concept was first set forth by Mendel in 1865 as a result of experiments
5 conducted on pea plants. Mendel proposed that a genetic determinant of a specific character is passed on from one generation to the next as a unit without any blending of the units, a theory now known as the "one gene one trait" theory.

Two of Mendel's proposed laws serve as the basis for modern
10 genetics. Mendel's first law states that two members (alleles) of a gene pair obligatorily segregate to different gametes, such that one half of the gametes carry one member (allele) of the pair and the other half of the gametes carry the other member (allele) of the gene pair. Mendel's second law states that during gamete formation the segregation of alleles
15 of a gene pair is independent of alleles other gene pairs.

Following Mendel's research, the discovery of chromosomes and the existence of linkage between genes closely positioned on a single chromosome have further contributed to our knowledge of inherited traits. Nowadays it is known that the second law of Mendel is valid for

genes residing on different chromosomes and for genes residing on the same chromosome, provided that they are at least 50 centiMorgan apart, whereas for closer genes residing on the same chromosome, linkage disequilibrium is experienced.

5 Thus, in an individual, a genetic trait in its simplest form is
determined by a pair of closely related sequences known as alleles. Each
allele, which is optionally an alternative form of a gene, resides on one of
the chromosomes of a chromosome pair. The pair of alleles of a given
individual can be identical in sequence or they can differ in sequence to
10 various degree. In meiosis, which is the process of cell division which
leads to the formation of sex cells (gametes), the chromosomes of any
given chromosome pair segregate into different gametes. This
chromosomal segregation is superseded by rearrangement of the DNA in
chromosome pairs which takes place via inter chromosomal
15 recombination events.

A specific trait of an individual is determined by a specific combination of two alleles carried by the individual. Thus, traits of an individual are determined by the genetic material inherited by that

individual from both parents as determined by chromosomal segregation and recombination events in each of the individual's parents.

Alleles carried on a chromosome pair encode the same genetic trait. Allelic variance ensures a high degree of distinction between
5 individuals.

Transformation of an exogene to a genome is typically random. Assuming one site of exogene integration, a resultant genome is said to be heterozygous of the type 1/0. By careful breeding and selection, a homozygous state for the exogene, i.e., 1/1 is obtainable. Although one
10 can transform a genome with a second exogene (2), it is practically impossible, using conventional transformation techniques, to direct the second exogene to a given integration site, in a different genome. As such, using such conventional molecular biology and breeding techniques, it is practically impossible to achieve exogene allelism of the
15 type 1/2.

Although it is theoretically possible to direct an exogene to a specific chromosomal site by gene knock-in techniques, which therefore could have been used to generate exogene allelism of the type 1/2, such

techniques, are very poor in transformation yields. Indeed, no attempts of generating exogene allelism were made so far.

The production of hybrid plants has been practiced ever since the beginning of this century. Plant breeders discovered that crossing two distinct parental lines often resulted in a hybrid plant which displays what is termed as hybrid vigor or heterosis, which is characterized by an increased crop yield and/or adaptation to both biotic and abiotic stresses. Early on, hybrid plant production concentrated mainly on corn and sorghum which were found to benefit from crossing of well compatible parental lines. Later it was discovered that hybrid plant production is also applicable to other plant species.

Several explanations have been proposed for the existence of heterosis. It is clear that a wider allelic variety exist in hybrids plant lines as is compared to inbred plant lines, since the alleles present in a hybrid plant are inherited from two distinct parent plant lines. This allelic variety is favored as a likely explanation for hybrid vigor, however, at present no scientific evidence has been brought forth to support this explanation.

Since hybrid plants have been demonstrated to be superior to inbred lines with respect to yield and vigor, the development of hybrid seeds is one of the prime objectives of the seed industry. In addition, since hybrid plant varieties result from a unique combination, the possibility of duplicating or reusing the hybrid seeds is minimized, thus, providing breeders with an inherent commercial protection.

The production of hybrid seed on a large scale is challenging because many crops have both male and female reproductive organs (stamen and pistil) on the same plant, either within a single flower or in separate flowers. This arrangement results in a high level of self pollination and makes large scale directed crosses between parental lines to generate hybrid plants difficult to accomplish.

To guarantee that out-crossing will occur during the production of hybrid seed, breeders have either manually or mechanically removed stamens from one parental line. Although such manual emasculation is effective for some plants, such as wheat, it is labor intensive and impractical for plants with small bi-sexual flowers. To cross-pollinate such plants breeders have often resorted to using naturally occurring male sterile mutants in efforts to produce hybrid seeds. Although the use of

naturally occurring male sterile mutants enables to cross pollinate plants with small bi-sexual flowers, the availability of such mutants and oftentimes the poor genetic makeup thereof severely limited wide spread use of this approach. In addition, using sterile mutants for out-crossing typically results in retention of sterility in a large fraction of the produced seeds. Such seeds when sown would then produce sterile plants which in the case of self pollinator crops such as wheat would lead to no crop yield for a fraction of the plants. To overcome the resultant low crop yield, a breeder is forced to select out the sterile seeds which can be a nearly impossible task.

Since important crops such as rice, wheat are self-pollinating plants with small bi-sexual flowers, there was a need and several attempts were made to develop systems for pollination control to assist in the production of F1 hybrids.

However, several conditions must exist in order to obtain economically feasible F1 hybrid plants. The male sterile line must be 100 % sterile and yet female fertile, a natural pollen transfer from the male fertile line to the male sterile line must be facilitated, and in cases of

grain or fruit crops, male fertility restoration (MFR) should be enabled in order to obtain crop yield in the F1 progeny.

There exist several mechanisms of male sterility in plants. One such mechanism is cytoplasmic male sterility (CMS). In general CMS has been used in corn, rice, sorghum and onion on a limited basis, see, for example, U.S. Pat. No. 3,861,079 to Patterson.

However, reliance on a single cytoplasmic-male-sterile system for the production of all hybrid plants is undesirable because it leaves the entire hybrid stock vulnerable to plant pathogens. For example, extensive use of one corn cytotyp, cmsT lead to an eptiphytic outbreak of Southern Corn Leaf Blight in the early 1970's. Thus, it is important to develop alternative methods to produce male sterile lines in plant species where only a single male-sterility system is available.

An emerging alternative to cytoplasmic sterility is a nuclear sterility. Nuclear male-sterile-based pollination systems rely upon the introduction of a male sterility trait to one parental plant followed by the introduction of a fertility-restorer gene, as a result of cross-pollination, to produce fertile hybrid plants.

Typically, genetically engineered nuclear male sterility is effected by expressing, in a controlled and targeted manner, a protein toxin which destroys anther tissues of the plant in which it is expressed. The expression of this toxin can be either silenced or antagonized in the F1 progeny by the introduction of a fertility-restorer gene, as a result of cross-pollination.

Male reproductive processes in flowering plants occur in the anther. This organ is composed of several tissues and cell types, and is responsible for producing pollen grains that contain the sperm cells. A specialized anther tissue, the tapetum, plays an important role in pollen formation. The tapetum surrounds the pollen sac early in anther development, degenerates during the later stages of development and is not present as an organized tissue in the mature anther. The tapetum produces a number of proteins and other substances that either aid in pollen development or become components of the pollen outer wall. It is known that many male sterility mutations interfere with tapetum cell differentiation and/or function. Thus tapetal tissue is believed to be essential for the production of functional pollen grains.

As such, many nuclear male-sterile-based pollination systems known in the art utilize anther specific promoters to target the destruction of the tapetum or other anther specific tissues during maturation such that subsequent pollen development is arrested.

5 For example, U.S. Pat. Nos. 5,409,823, 5,659,124 and 5,824,542 to Crossland describe a dual method for producing male-sterile plants. Two genetically transformed plants, parents 1 and 2, are crossed to obtain male-sterile offspring. Parent 1 is transformed with an expression cassette comprising a nucleotide sequence encoding an anther-specific
10 promoter which is operably linked to a nucleotide sequence encoding a trans-activator. Parent 2 is transformed with an expression cassette comprising a target nucleotide sequence, which is capable of being activated by the trans-activator, operably linked to a nucleotide sequence which encodes a toxin which can be an RNA or a polypeptide and which
15 disrupts the formation of viable pollen. Therefore, crossing parent 1 with parent 2 results in male-sterile offspring. The male-sterile plants produced are useful for producing hybrid seed. Subsequent crossing of the male sterile plant with a plant which does not include the trans-activator restores fertility.

Although this system produces desirable male sterility in a portion of the produced offspring, restoration of fertility which relies upon the segregation of the trans-activator gene from the toxin gene cannot be efficiently effected using this system. In the male sterile plant the segregation of these two genes during meiosis relies a great deal on their physical chromosomal location, as such when a male sterile plant is crossed with a plant which does not express the trans-activator, the trans-activator and the toxin can be provided by the male sterile parent as a result of which a portion of the produced progeny will retain undesirable male sterility.

Thus the method by Crossland is not efficiently applicable for self pollinators such as, for example, wheat, cotton and rice.

U.S. Pat. No. 5,929,307 to Hodges describes a recombinant expression vector comprising a suicide gene flanked by site-specific recombination sequences, which vector, when introduced into a plant leads to male sterility. This invention further relates to a second expression vector including a recombinase gene which when introduced into the male sterile plant via a cross-pollinating restorer plant, leads to the deletion of the toxin gene and as a result to the restoration of fertility.

It will be appreciated in this case that since restoration of fertility is effected by a gene product which is introduced via a cross-pollinating restorer plant, a high copy number of the restorer gene must be maintained in the restorer plant such that highly efficient restoration of fertility in the resultant progeny is effected.

Thus, the presently available nuclear male-sterility pollination systems, although generally efficient in the production of male sterile plants, suffer from limitations when restoration of such male sterility via out-crossing is desirable, such as the case for grain or fruit producing hybrid plants. Since these systems cannot enable 100 % restoration of fertility to the male sterile plants produced and since the resultant male sterile hybrid seeds are not typically selected out, a suboptimal crop yield from grain or fruit producing hybrid plants produced by these systems results.

There is thus a widely recognized need for, and it would be highly advantageous to have, a male sterile plant, which plant when out-crossed with a compatible male fertile plant generates male fertile offspring in substantially 100 % of it's progeny. As further detailed hereinunder, such a plant can be generated by a transformation method which results

in formation of allelism between two exogenes. There is thus a widely recognized need for, and it would be highly advantageous to have, a method and expression cassettes for the generation of exogene allelism.

5

SUMMARY OF THE INVENTION

It is one object of the present invention to provide a eukaryotic organism, such as an organism of the *animalia* or *planta* kingdom, or a commercially important product thereof, having a genome characterized
10 by exogenic allelism, i.e., allelism of two distinct exogenes, which can be used to determine a phenotype of the organism.

It is another object of the present invention to provide expression cassettes which can be used by preplanned transformation and crossing regime to generate a eukaryotic organism having a genome characterized
15 by exogenic allelism.

It is yet another object of the present invention to provide methods of generating exogenic allelism in a eukaryotic organism exploiting the expression cassettes of the former object of the invention.

It is still another object of the present invention to exploit the above methods and expression cassettes for the generation of substantially 100 % reversible plant male sterility, exploitable, for example, in the hybrid seed industry of grain and fruit crops characterized
5 by small, bi-sexual flowers.

Thus, according to one aspect of the present invention there is provided a non-human eukaryotic organism having a genome comprising a first exogene being in a first chromosome of a chromosome pair of the genome and a second exogene being in a second chromosome of the
10 chromosome pair, the first and the second exogenes being in allelic relationship, such that the first and the second exogenes obligatorily segregate to different gametes.

According to further features in preferred embodiments of the invention described below, expression of the first and the second
15 exogenes determines a phenotype of the organism.

According to still further features in the described preferred embodiments an expression product of the second exogene transactivates expression of the first exogene.

According to still further features in the described preembodiments the second exogene encodes an RNA polymerase.

According to still further features in the described preferred embodiments the second exogene encodes a transcription factor.

5 According to still further features in the described preferred embodiments the first exogene encodes a polypeptide selected from the group consisting of a cytotoxic polypeptide and a cytostatic polypeptide.

According to still further features in the described preferred embodiments the first exogene encodes an RNA molecule selected from
10 the group consisting of an antisense RNA molecule and a ribozyme RNA molecule.

According to still further features in the described preferred embodiments expression products of the first and the second exogenes assemble into a hetero-oligomeric protein.

15 According to still further features in the described preferred embodiments the hetero-oligomeric protein has an activity selected from the group consisting of cytotoxic activity and cytostatic activity.

According to still further features in the described preferred embodiments the organism is a plant species.

According to still further features in the described preferred
embodiments the second exogene encodes an RNA polymerase non-
operable with eukaryotic promoters.

According to still further features in the described preferred
5 embodiments the RNA polymerase is selected from the group consisting
of bacterial RNA polymerase and bacteriophage RNA polymerase.

According to still further features in the described preferred
embodiments the bacteriophage RNA polymerase is selected from the
group consisting of T7 RNA polymerase, T3 RNA polymerase and SP6
10 RNA polymerase.

According to still further features in the described preferred
embodiments the first exogene encodes a polypeptide selected from the
group consisting of a cytotoxic polypeptide and a cytostatic polypeptide.

According to still further features in the described preferred
15 embodiments the polypeptide is selected from the group consisting of
pectate lyase, 1-3 β -glucanase, avidin, streptavidin, diphtheria Toxin A-
chain (DTA), URF13, Indole acetic acid-lysine synthetase, CytA toxin
and RNase-TI.

According to still further features in the described preferred
embodiments the specific tissue forms a part of an stamen tissue of the
plant.

According to another aspect of the present invention there is provided an expression cassette comprising (a) a first segment including a first transcribable polynucleotide sequence, the first transcribable polynucleotide sequence being operatively linked to a first promoter sequence, the first segment being flanked by a pair of first site-specific recombination sequences; and (b) a second segment, being linked to the first segment, the second segment including a second transcribable polynucleotide sequence, the second transcribable polynucleotide sequence being operatively linked to a second promoter sequence, the

second segment being flanked by a pair of second site-specific recombination sequences.

According to yet another aspect of the present invention there is provided a method of generating exogenic allelism in a non-human eukaryotic organism, the method comprising the steps of (a) generating a first and second isogenic organisms homozygous for the expression cassette described hereinabove; (b) introducing a first recombinase into the first organism, so as to excise the first segment; (c) introducing a second recombinase into the second organism, so as to excise the second segment; and (d) crossing the organisms resultant from steps (b) and (c), so as to generate an offspring characterized by exogenic allelism.

According to further features in preferred embodiments of the invention described below, the first and the second pair of directly repeated site-specific recombination sequences are each independently selected from the group consisting of Lox recombination sequences, FRT recombination sequences, Gin recombinase sequences, Pin recombinase sequences and R/RS recombinase sequences.

According to still another aspect of the present invention there is provided an expression cassette comprising (a) a first segment including

a first promoter sequence; (b) a second segment including a first transcribable polynucleotide sequence; and (c) a third segment including a second transcribable polynucleotide sequence, the second transcribable polynucleotide sequence being operatively linked to a second promoter sequence, the third segment being flanked by the first and second segments, wherein a pair of site-specific recombination sequences are disposed one between the first segment and the third segment and another between the second segment and the third segment, such that the first promoter sequence is operatively coupled with the first transcribable polynucleotide sequence only following excision of the third segment from the expression cassette by site specific recombination via the pair of site-specific recombination sequences.

According to an additional aspect of the present invention there is provided a method of generating exogenic allelism in a non-human eukaryotic organism, the method comprising the steps of (a) generating a first and a second isogenic organisms homozygous for the expression cassette described hereinabove; (b) introducing a recombinase into the first organism, so as to excise the third segment thereby operatively adjoining the first transcribable polynucleotide sequence to the first

promoter sequence; and (c) crossing the organism resultant from step (b) and the second organism, so as to generate an offspring characterized by exogenic allelism.

According to further features in preferred embodiments of the invention described below, the first promoter sequence is a non-eukaryotic promoter sequence.

According to still further features in the described preferred embodiments the second promoter sequence is a tissue specific promoter sequence naturally operable in a first tissue.

According to still further features in the described preferred embodiments the first promoter sequence is a tissue specific promoter sequence naturally operable in a second tissue and further wherein the second transcribable polynucleotide sequence encodes a tissue specific transcription activator naturally expressed in the second tissue and naturally capable of activating the first promoter sequence.

According to still further features in the described preferred embodiments the second transcribable polynucleotide sequence encodes a transactivator.

According to still further features in the described preferred embodiments the transactivator is an RNA polymerase.

According to still further features in the described preferred embodiments the first transcribable polynucleotide sequence encodes an
5 enzyme.

According to still further features in the described preferred embodiments the first transcribable polynucleotide sequence encodes an RNA molecule selected from the group consisting of an antisense RNA molecule and a ribozyme RNA molecule.

10 According to still further features in the described preferred embodiments the second transcribable polynucleotide sequence encodes a transactivator of the first promoter sequence.

According to yet an additional aspect of the present invention there is provided a plant homozygous for any of the expression cassettes
15 herein described.

According to still an additional aspect of the present invention there is provided a plant comprising a genome, the genome including a pair of exogenes being in allelic relationship, wherein a first exogene of the pair of exogenes being located on a first chromosome of a

chromosome pair of the genome of the plant, and further wherein a second exogene of the pair exogenes being located on a second chromosome of the chromosome pair of the genome of the plant.

According to further features in preferred embodiments of the invention described below, the first and second exogenes are selected such that expression thereof generates male sterile plant.

According to still further features in the described preferred embodiments by crossing the male sterile plant with a male fertile plant results in offsprings characterized by male fertility.

According to a further aspect of the present invention there are provided plant seeds each of which comprising a genome, the genome including a pair of exogenes being in allelic relationship, wherein a first exogene of the pair of exogenes being located on a first chromosome of a chromosome pair of the genome of the plant seeds, and further wherein a second exogene of the pair of exogenes being located on a second chromosome of the chromosome pair of the genome of the plant seeds.

The present invention successfully addresses the shortcomings of the presently known configurations by providing a eukaryotic organism having a genome characterized by exogenic allelism, i.e., allelism of two

distinct exogenes, which can be used to determine a phenotype of the organism, expression cassettes and methods useful in the generation of such an organism.

5

BRIEF DESCRIPTION OF THE DRAWINGS

The invention is herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the cause of providing what is believed to be the most useful and readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention in more detail than is necessary for a fundamental understanding of the invention, the description taken with the drawings making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

In the drawings:

FIG. 1 is a flow diagram outlining one method for generating plants according to the teachings of the present invention. TapP - tapetum promoter; Lox - Cre recombinase recognition site; T7P - bacteriophage T7 promoter; T7 polymerase - the bacteriophage RNA
 5 polymerase recognizes the T7 promoter site; 35S - the CaMV promoter; CRE - the CRE recombinase.

FIG. 2 is a flow diagram outlining another method for generating plants according to the teachings of the present invention. TapP - tapetum promoter; Lox - Cre recombinase recognition site; T7P -
 10 bacteriophage T7 promoter; T7 polymerase - the bacteriophage RNA polymerase recognizes the T7 promoter site; 35S - the CaMV promoter; CRE - the CRE recombinase; FRT - the FLP recombinase recognition site.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention reduces to practice a novel concept in genetic engineering, according to which a eukaryotic organism is transformed and crossed so as to provide an offspring including a first
5 exogene in a first chromosome of a chromosome pair and a second exogene in a second chromosome of said chromosome pair, the exogenes exhibiting allelic relationship, such that the exogenes obligatorily segregate to different gametes. The present invention is further of expression cassettes for implementing the novel concept, methods of
10 implementing same for the obtainment of reversible male sterility in plants, and plants and plant products obtained thereby.

The principles and operation of a methods and cassettes for generating exogenic allelism in a non-human eukaryotic organism according to the present invention may be better understood with
15 reference to the drawings and accompanying descriptions.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the

drawings. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

5 According to one aspect of the present invention there is provided a non-human eukaryotic organism having a genome comprising a first exogene in a first chromosome of a chromosome pair of the genome and a second exogene in a second chromosome of the chromosome pair. The first and second exogenes exhibiting allelic relationship, such that they
10 obligatorily segregate to different gametes when such gametes form in the organism.

 It will be appreciated that since non-human eukaryotic organisms contain a plurality of chromosome pairs, an allelic relationship of two exogenes can be established on more than one chromosome pair and in
15 more than one location on a given chromosome pair .

 As used herein the phrase "allelic relationship" and the term "allelism" refer to the positional relationship between two genes on two chromosomes of a chromosome pair. As such "exogenic allelism" is the allelic positioning of two functionally distinct exogenes on the

chromosomes of a chromosome pair such that substantially 100 % segregation of the two exogenes is observed upon gamete formation.

As used herein the term "exogenes" refers to polynucleotide sequences which are trans-introduced into and integrated in a genome of a species. Such exogenes may be of a different or the same species. Exogenes according to the present invention include transcribable sequences, e.g., mRNA and protein coding sequences, both cDNA and genomic sequences, and antisense RNA coding sequences, and/or cis acting sequences which cis-control gene expression, such as promoters and enhancers. Such exogenes, assembled in an expression cassette, are used to transform cells and/or organisms to thereby obtain transformed, genetically modified or transgenic cells and/or organisms harboring the exogene(s) in their genome. A transgenic organism is a stably transformed organism having one or more cells that contain an exogene. The term stably transformed refers to a transformed cell or organism (host) that is capable of transmitting an exogene to its progeny. According to the present invention a stably transformed host has an exogene integrated into its genome. Exogenes according to the present invention may represent sequences which naturally occur in the

organism, or mutants, portions and chimeras thereof or sequences which do not naturally occur in the organism, for example sequences from a different sub species, species or genus.

According to a preferred embodiment of the present invention the
 5 expression of the first and the second exogenes determines a phenotype of the organism. As used herein the term "phenotype" refers to characteristic gene and/or exogene expression which results in the presence of distinct molecular product(s) in an organism. The presence of such molecular product(s) such as for example a protein or an RNA
 10 molecule (e.g. mRNA or antisense RNA) may lead to morphological changes, differential resistivity to pathogens and chemicals, differentiated development and the like.

Thus, the expressed exogenes co-affect the phenotype of an organism, that is to say that the presence of expression products from
 15 both exogenes is required in order to achieve a certain phenotype.

The term "expression" refers to the biosynthesis of a gene product. For example, in the case of a structural gene, expression involves the transcription of the structural gene into messenger RNA (mRNA) and the

translation of the mRNA into one or more polypeptides. In the case of antisense RNA or ribozymes the expression product is an RNA molecule.

A trait determined by two allelic exogenes is particularly advantageous and is the gist of the present invention since crossing a

~~transgenic org characterized by exogenic allelism with a second organism~~

leads to the complete segregation of the two exogenes and as such to loss

of the phenotype in the offsprings. This cannot be achieved for traits

determined by a single gene or even a multiplicity of genes arranged in a

non-allelic fashion, since complete segregation of these genes and as

10 such loss of the trait cannot be ensured. The advantages of exogenic

allelism are further exemplified hereinbelow with respect to plants

characterized by exogenic allelism.

According to one preferred embodiment of the present invention

an expression product of the second exogene transactivates expression of

15 the first exogene.

The transactivators of the present invention include, but are not

limited to, polymerases, DNA binding proteins, naturally occurring and

synthetic transcriptional activators, translational activators, post-

transcriptional activators, and the like. It will be appreciated that only

transactivators which do not naturally occur in the cells or tissue in which they are to be expressed are selected. Preferably, transactivators from unrelated organisms are selected.

According to one preferred embodiment of the present invention

5 the transactivator is a bacteriophage RNA polymerase. The use of such transactivator polypeptides in directing expression of another nucleotide sequence is exemplified by the T7 RNA polymerase. See, U.S. Pat. Nos. 5,122,457; 5,126,251; and 5,135,855; Lassner *et al.* (1991) Plant Molecular Biology 17:229-2; Rodriguez *et al.* (1990) Journal of Virology

10 64:4851-4857; Vennema *et al.* (1991) Gene 108:201-210; Benton *et al.*, Molecular and Cellular Biology (1990) Molecular and Cellular Biology 10:353-360; Elroy-Stein and Moss (1990) Proceedings, Proc. Natl. Acad. Sci.:USA 87:6743-6747; Moss *et al.* (1990) Nature 348:91-92; Elroy-Stein *et al.* (1989) Proceedings, Proc. Natl. Acad. Sci.:USA 86:6126-

15 6130; and Rosenberg *et al.* (1987) Gene 56:125-135, herein incorporated by reference. It will be appreciated that other polymerases which do not normally exist in the organism such as, for example, polymerases from species different than that of the organism but yet of the same genus, can also be utilized by the present invention.

Transactivators also include transcriptional activators which are necessary for transcription activation of specific promoters. Binding domains of one protein may be fused to activity domains of another protein to form chimeras of such DNA binding proteins, such as

5 GAL4/VP16 (Carey *et al.* (1989), J. Mol. Biol., 209:423-432; Cress *et al.* (1991) Science, 251:87-90; and Sadowski *et al.* (1988), Nature, 335:563-564). Likewise, the binding domain of other proteins, i.e., Lex A (Brent and Ptashne, (1985), Cell, 43:729-736, which describes a Lex A/GAL4 transcriptional activator) can be utilized.

10 Transactivators can also be translational activators. Translational activators are exemplified by the cauliflower mosaic virus translational activator (TAV). See, for example Futterer and Hohn (1991) EMBO J. 10:3887-3896. In this system a dicistronic mRNA is produced. That is, two coding regions are transcribed in the same mRNA from the same

15 promoter. In the absence of TAV, only the first cistron is translated by the ribosomes. However, in cells expressing TAV, both cistrons are translated.

Thus, when expressed, the transactivator transactivates the expression of the first exogene. Thus, it is the expression of the first

exogene which determines the phenotype of the organism, but since the expression of the first exogene is dependent on the presence of the second exogene product, both must be expressed in order to establish and maintain a phenotype.

5 According to preferred embodiments of the present invention the first exogene encodes a molecule which can either arrest cell proliferation and/or cause cell degeneration.

6649374
10 Thus, according to one preferred embodiment of the present invention the first exogene encodes a cytotoxic polypeptide or a cytostatic polypeptide. The expression of the first exogene either prevents the cell in which it is expressed from proliferating or alternatively it causes cell degeneration and ultimately cell death. Specific examples of cytostatic and cytotoxic polypeptides are further described hereinbelow.

15 In an alternative embodiment, the first exogene encodes an antisense RNA or a ribozyme which when transcribed target specific mRNA sequences expressed within the cell and as such can be used, for example, to disrupt biological processes and bring about cell death.

According to another preferred embodiment of the present invention the expression products of the first and the second exogenes assemble into a hetero-oligomeric protein. In this case both exogenes are preferably concomitantly expressed such that accumulation of the two non-functional expression products leads to assembly of a functional hetero-oligomer.

The first and second exogenes are also referred to hereinbelow as the first and second transcribable polynucleotide sequences.

According to another aspect of the present invention there is provided an expression cassette useful in generating exogenic allelism in a non-human eukaryotic organism. The expression cassette according to this aspect of the present invention includes a first segment including a first promoter sequence. The expression cassette further includes a second segment including a first transcribable polynucleotide sequence and a third segment flanked by the first and second segments and including a second transcribable polynucleotide sequence operatively linked to a second promoter sequence. A pair of site-specific recombination sequences are disposed one between the first segment and the third segment and another between the second segment and the third

segment. Thus, the first promoter sequence can be operatively coupled with the first transcribable polynucleotide sequence following excision of the third segment from the expression cassette by site specific recombination via the pair of site-specific recombination sequences.

5 As used herein an expression cassette is a polynucleotide molecule comprising at least one polynucleotide sequence that is expressed in a host cell or organism. Typically such expression is under the control of certain cis acting regulatory elements including constitutive, inducible or tissue-specific promoters, and enhancer elements. Common to the art, 10 such polynucleotide sequence(s) are said to be "operably linked to" the regulatory elements. Expression cassettes typically also include eukaryotic or bacterial derived selectable markers that allow for selection of eukaryotic cells containing the expression cassette. These can include, but are not limited to, various genes which confer antibiotic resistance 15 and which are well known in the art and therefore will not be further described herein.

For many applications it is required that the expression cassette described herein will be integrated in a nucleic acid construct, such as an expression construct or an antisense construct. Such constructs are well

known in the art, are commercially available and may include additional sequences, such as, for example, a cloning site, one or more prokaryote or eukaryote marker genes with their associated promoters for selection of prokaryotic cells containing the expression cassette, one or more
5 prokaryotic origins of replication, one or more translation start site, one or more polyadenylation signal, and the like.

A construct according to the present invention preferably further includes an appropriate selectable marker. In a more preferred embodiment according to the present invention the construct further
10 includes an origin of replication. In another most preferred embodiment according to the present invention the construct is a shuttle vector, which can propagate both in *E. coli* (wherein the construct comprises an appropriate selectable marker and origin of replication) and be compatible for propagation in cells, or integration in the genome, of an
15 organism of choice. The construct according to this aspect of the present invention can be, example, a plasmid, a bacmid, a phagemid, a cosmid, a phage, a virus or an artificial chromosome.

To be expressed, polynucleotide sequences included in an expression cassettes must be driven by a promoter. Several types of

promoters are now well known in the art of transformation, as are other regulatory elements that can be used alone or in combination with such promoters.

As used herein the term "promoter" includes reference to a region
5 of DNA upstream from the start of transcription and involved in recognition and binding of RNA polymerase and other proteins which are required to initiate transcription. Promoters which initiate transcription only in certain tissues are referred to as "tissue-specific". A "cell type" specific promoter primarily drives expression in certain cell types in one
10 or more organs.

According to a preferred embodiment of the present invention the second promoter of the expression cassettes described hereinabove, and which drives the expression of the transactivator is a tissue specific promoter. It will be appreciated, however, that other types of promoters
15 such as inducible promoters, constitutive promoters and developmentally regulated promoters can also be used by the present invention depending on the desired spatial and temporal expression of the phenotype determined by the exogenic allelism.

Specific examples to polynucleotide sequences and promoters are detailed hereinbelow with respect to the plant embodiments of the present invention which are further exemplified in Examples 1 and 2 of the Examples section.

5 A number of different site-specific recombinase systems can be utilized in accordance with the present invention, including, but not limited to, the Cre/lox system of bacteriophage P1, the FLP/FRT system of yeast, the Gin recombinase of phage Mu, the Pin recombinase of *E. coli*, and the R/RS system of the pSR1 plasmid. The presently two
10 preferred site-specific recombinase systems are the bacteriophage P1 Cre/lox and the yeast FLP/FRT systems, for these systems have shown to have high recombination yields by others. The site specific recombination sequence for each of these two systems is relatively short (34 bp for lox and 34-47 bp for FRT). In these systems a recombinase
15 (Cre or FLP) interacts specifically with its respective site-specific recombination sequence (lox or FRT, respectively) to invert or excise the intervening sequences depending on the orientation of the site-specific recombination sequences. The site specific recombination sequences utilized by the expression cassettes of the present invention are oriented

as direct repeats, so as to affect excision of the sequence flanked thereby.

Site-specific recombination systems are further described in U.S. Pat. No. 5,527,695, which is incorporated herein by reference.

As mentioned hereinabove, to effect excision a suitable
 5 recombinase is introduced into the target organism by any suitable
 method which leads to the expression of the recombinase within the
 organism. For example, the recombinase gene may be stably integrated
 into the genome of the organism or alternatively transiently expressed
 therein. As further detailed hereinbelow with respect to plants, a
 10 recombinase can also be introduced into a plant via sexual crossing with
 a transgenic plant carrying the recombinase gene. Similar crossings can
 be made between, for example, inbred organisms of the animalia
 kingdom.

To generate exogenic allelism in the organism, the above
 15 described expression cassette is introduced into the genome of the
 organism by transformation methods known in the art, some of which are
 described in detail below.

Following establishment of a stably transformed organism, a first
 and a second isogenic organisms homozygous for the expression cassette

are propagated therefrom. A recombinase gene specific for the site specific recombinase sequences is introduced into the first organism, so as to excise the third segment thereby operatively adjoining the first transcribable polynucleotide sequence to the first promoter sequence. As
 5 already mentioned, the recombinase gene is either introduced via transformation techniques or preferably via out-crossing with a recombinase expressing organism which does not contain the expression cassette, but which is preferably isogenic to the first organism.

Finally the resultant first and second organisms are crossed so as
 10 to generate an offspring characterized by exogenic allelism. Since each organism contributes a single chromosome of a chromosome pair to it's offspring, such an offspring will contain as a part of it's genome a first expressible polynucleotide on a first chromosome of a chromosome pair and a second expressible polynucleotide on a second chromosome of the
 15 chromosome pair, wherein the first and second expressible polynucleotides exhibit allelism, i.e., such polynucleotides obligatorily segregate to different gametes.

It will be appreciated that when a single pair of site specific sequences is used, the expression cassette positioning of the first

polynucleotide coding sequence and the second polynucleotide coding sequence can be switched around.

As is further exemplified hereinabove the first transcribable polynucleotide sequence encodes a cytotoxic or cytostatic molecule and
5 the second polynucleotide encodes a transactivator. In this case the expression of the transactivator will be dependent on the type of the second promoter used and will most typically be, in cases wherein a constitutive or tissue specific promoter is used, initiated following transformation of the expression cassette into the organism. On the other
10 hand, in an alternative polynucleotide sequences arrangement of this expression cassette, the transactivator can be encoded by the first transcribable polynucleotide sequence, while the cytotoxic or cytostatic molecule can be encoded by the second transcribable polynucleotide sequence. Thus, in this case the transactivator will be expressed only
15 following recombination. As such, in the latter configuration, which is described in more detail hereinbelow in Example 1 of the examples section, expression from any exogenic sequences will initiate only following recombination. This is particularly advantageous in situations

wherein the transactivator is toxic to the cell in which it is expressed. In such cases premature expression of the transactivator must be avoided.

As is further detailed hereinunder, exogenic allelism can also be generated using an expression cassette which utilizes two pairs of site specific recombination sequences.

Thus, according to another aspect of the present invention there is provided an expression cassette useful in generating exogenic allelism in a non-human eukaryotic organism. The expression cassette according to this aspect of the present invention includes a first segment flanked by a pair of first site-specific recombination sequences and including a first transcribable polynucleotide sequence, wherein the first transcribable polynucleotide sequence is operatively linked to a first promoter sequence.

The expression cassette according to this aspect of the present invention further includes a second segment, linked to the first segment, the second segment including a second transcribable polynucleotide sequence operatively linked to a second promoter sequence, wherein the second segment is flanked by a pair of second site-specific recombination sequences.

Following establishment of a stably transformed organism using the above cassette via transformation as is further detailed hereinbelow, a first and a second isogenic organisms, homozygous for the expression cassette, are propagated therefrom. A first recombinase gene is introduced, as described above, into the first organism, so as to excise the first segment. A second recombinase gene is introduced, as described above, into the second organism, so as to excise the second segment.

Finally the resultant first and second organisms are crossed so as to generate an offspring characterized by exogenic allelism.

As before, since each organism contributes a single chromosome of a chromosome pair to its offspring, such an offspring will contain as a part of its genome a first expressible polynucleotide on a first chromosome of a chromosome pair and a second expressible polynucleotide on a second chromosome of the chromosome pair.

Further related description of the expression cassette according to this aspect of the present invention is given hereinbelow in Example 2.

According to yet another aspect of the present invention and as demonstrated in Examples 1 and 2 of the examples section exogenic allelism is used to generate reversible nuclear male sterility in plants.

Male sterility is the failure or inability to produce functional or viable pollen. Male sterility may result from defects leading to the non-formation of pollen or to the lack of functional ability in the pollen when it is formed. Therefore, either pollen is not formed or, if formed, it is
5 either non-viable or incapable of effective fertilization under normal conditions.

The male-sterile plants of the present invention, are female fertile. That is, the plants do not produce fertile pollen, yet are capable of accepting pollen from the desired paternal parent resulting in fertilization
10 and hybrid seed production.

To effect male sterility in a plant, the two transcribable polynucleotide sequences must be expressed such that when expressed cytostatic or cytotoxic activity only occurs in the stamen tissue or tissues (e.g., anther, pollen or tapetum).

15 Thus, according to a preferred embodiment of this aspect of the present invention, the second promoter is an stamen tissue-specific promoter. As used herein the term "stamen" refers to the male fertilizing organ of a flowering plant, including anther tissues. As used herein the

phrase "anther tissues" or the term "anther" is meant to include the pollen and tapetum tissues as well.

In the case of promoter DNA sequences, stamen specific promoters are typically specific to the anther tissues and include
5 regulatory sequences which direct the transcription of associated transcribable sequences so that the corresponding RNA is present in anther tissues in concentrations at least 100-fold that observed in other tissues.

Anther-specific promoters are well known in the art, examples of
10 which include, but are not limited to, a tapetum-specific promoter such as the tobacco anther promoter, *ant32*, an anther-specific promoter such as that from LAT52 (Twell *et al.*, Mol. Gen. Genet. 217: 240-245 (1989)); a pollen-specific promoter such as that from Zm13 (Guerrero *et al.*, Mol. Gen. Genet. 224: 161-168 (1993)) or a microspore-preferred promoter
15 such as that from apg (Twell *et al.*, Sex. Plant Reprod. 6: 217-224 (1993). Additional examples are provided in, for example, U.S. Pat. No. 5,477,002, which discloses promoter sequences from anther-specific genomic clones which are operatively linked to a DNA sequence coding for a desired polypeptide to enable anther specific expression.

By utilizing stamen-specific promoters, the resulting transgenic plants will express the transactivator only in the stamen tissues of the plant.

As already mentioned hereinabove, a transactivator expressed
5 from the second transcribable polynucleotide sequence of an expression cassette can be an RNA polymerase, a transcriptional activator or a translational activator.

In a preferred embodiment of this aspect of the present invention the transactivator is a bacteriophage RNA polymerase such as, but not
10 limited to, a T7 RNA polymerase, a T3 RNA polymerase or an SP6 RNA polymerase.

As such the first promoter which drives the expression of the first transcribable polynucleotide sequence includes a sequence recognizable by bacteriophage RNA polymerase such that in the presence of the
15 bacteriophage RNA polymerase and providing operational linkage between the first transcribable polynucleotide sequence and the first promoter, transcription of the first polynucleotide sequence is effected.

According to a preferred embodiment of this aspect of the invention the expression product of the first transcribable polynucleotide sequence is utilized to disrupt formation of viable pollen when expressed.

As mentioned hereinabove the resultant expression product of the first transcribable polynucleotide sequence can be either a polypeptide or an RNA molecule such as for example an anti-sense RNA molecule or a ribozyme RNA molecule.

Examples of polypeptides suitable for use in this aspect of the present invention include, but are not limited to, proteins capable of inhibiting the synthesis of macromolecules that are essential for cellular function, enzymes that degrade macromolecules that are essential for cellular function, proteins that alter the biosynthesis or metabolic metabolism of plant hormones and proteins that inhibit a specific function or development of anther/tapetum cells.

For example, Mariani *et al.*, Nature, 347:737, (1990), have shown that expression of either *Aspergillus oryzae* RNase-TI or an RNase of *Bacillus amyloliquefaciens*, designated "BARNASE", in the tapetal cells of a plant induced destruction of the tapetal cells, resulting in male sterility. Other genes can be used as alternatives to BARNASE for the

development of male sterile plants, such as an anther-specific β -1,3-glucanase (Hird *et al.* The Plant Journal 4:1023-1033, 1993), or the male sterility gene described by Aarts *et al.* Nature 363: 715-717, 1993.

Additional polypeptides include diphtheria Toxin A-chain (DTA),
 5 which inhibits protein synthesis, Greenfield *et al.* (1983), Proc. Natl. Acad., Sci.:USA, 80:6853; Palmiter *et al.* (1987), Cell, 50:435; Pectate lyase pelE from *Erwinia chrysanthemi* EC16, which degrades pectin, causing cell lysis. Keen *et al.* (1986), J. Bacteriology, 168:595; T-urf13 (TURF-13) from cms-T maize mitochondrial genomes; this gene encodes
 10 a polypeptide designated URF13 which disrupts mitochondrial or plasma membranes. Braun *et al.* (1990), Plant Cell, 2:153; Dewey *et al.* (1987), Proc. Natl. Acad. Sci.:USA, 84:5374; Dewey *et al.* (1986), Cell, 44:439; Gin recombinase from phage Mu a gene, which encodes a site-specific DNA recombinase which will cause genome rearrangements and loss of
 15 cell viability when expressed in cells of plants. Maeser *et al.* (1991), Mol. Gen. Genet., 230:170-176; Indole acetic acid-lysine synthetase (iaaL) from *Pseudomonas syringae*, which encodes an enzyme that conjugates lysine to indole acetic acid (IAA). When expressed in the cells of plants, it causes altered developments due to the removal of IAA from the cell

via conjugation. Romano *et al.* (1991), Genes and Development, 5:438-446; Spena *et al.*, Mol. Gen. Genet., (1991), 227:205-212; Roberto *et al.* (1992 XXX) Proc. Natl. Acad. Sci.:USA, 87:5795-5801; CytA toxin gene from *Bacillus thuringiensis Israeliensis* which encodes a protein that is

5 mosquitocidal and hemolytic. When expressed in plant cells, it causes death of the cell due to disruption of the cell membrane. McLean *et al.* (1987), J. Bacteriology, 169:1017-1023; Ellar *et al.* (1990), U.S. Pat. No. 4,918,006; and, biotin binding proteins such as Streptavidin and avidin as further detailed in WO 96/40949 and 99/04023 which describe the use of

10 a biotin binding protein for the induction of male sterility.

In addition, the first polypeptide sequence can also encode DNase, RNase; protease; salicylate hydroxylase and the like.

In cases where it may be beneficial to target the polypeptide product expressed from the first transcribable polynucleotide sequence to

15 a subcellular compartment such as the chloroplast, vacuole, peroxisome, glyoxysome, cell wall or mitochondrion, or for secretion into the apoplast, the first transcribable polynucleotide sequence includes a signal sequence, 5' and/or 3' to the region of the polypeptide encoding sequence. Targeting sequences at the 5' and/or 3' end of the structural gene may

determine, during protein synthesis and processing, where the encoded protein is ultimately compartmentalized. The presence of a signal sequence directs a polypeptide to either an intracellular organelle or subcellular compartment or for secretion to the apoplast. Many

5 sigsequences are known in the art. To this end, see, for example, Becker *et al.*, Plant Mol. Biol. 20: 49 (1992), Close, P. S., Master's Thesis, Iowa State University (1993), Knox, C., *et al.*, "Structure and Organization of Two Divergent Alpha-Amylase Genes From Barley", Plant Mol. Biol. 9: 3-17 (1987), Lerner *et al.*, Plant Physiol. 91: 124-129 (1989), Fontes *et*

10 *al.*, Plant Cell 3: 483-496 (1991), Matsuoka *et al.*, Proc. Natl. Acad. Sci. 88: 834 (1991), Gould *et al.*, J. Cell Biol 108: 1657 (1989), Creissen *et al.*, Plant J. 2: 129 (1991), Kalderon, D., Robers, B., Richardson, W., and Smith A., "A short amino acid sequence able to specify nuclear location", Cell 39: 499-509 (1984), Stiefel, V., Ruiz-Avila, L., Raz R., Valles M.,

15 Gomez J., Pages M., Martinez-Izquierdo J., Ludevid M., Landale J., Nelson T., and Puigdomenech P., "Expression of a maize cell wall hydroxyproline-rich glycoprotein gene in early leaf and root vascular differentiation", Plant Cell 2: 785-793 (1990).

Alternatively male sterility can be induced by the expression of an antisense RNA. The binding of an antisense messenger RNA molecule to target messenger RNA (mRNA) molecules results in RNaseH associated degradation of the mRNA and/or arrest of translation. Thus, a
5 suitable antisense molecule would have a sequence that is complementary to that of a messenger RNA species encoding a protein that is essential for tapetal cell function or pollen synthesis.

For example, an antisense RNA that interferes with the expression of the RTS2 gene product can be utilized by the present invention.

10 Alternatively, the antisense of PRK1 (pollen-expressed receptor-like kinase) from *Petunia inflata* (Mu *et al.* The Plant Cell 6:709-721, 1994), or the antisense directed against the Bcp1 male fertility gene of *Arabidopsis* (Xu *et al.* Proc. Natl. Acad. Sci. 92: 2106-2110, 1995) can also be utilized as an antisense RNA according to the present invention.

15 In addition an antisense RNA molecule can be directed to a nuclear localization signal sequence and as such bring about degradation and/or arrest of translation of mRNA species which contain NLS signal sequences.

The first polynucleotide can also encode a ribozyme. Ribozymes can be designed to express endonuclease activity that is directed to a certain target sequence in a messenger RNA molecule. For example, Steinecke *et al.*, EMBO J., 11:1525, (1992), achieved up to 100 % inhibition of neomycin phosphotransferase gene expression by ribozymes in tobacco protoplasts. Alternatively, an RNA molecule capable of promoting RNaseP-mediated cleavage of target messenger RNA molecules can also be used. According to this approach, an external guide sequence can be constructed for directing the endogenous RNaseP to a particular species of intracellular messenger RNA which is subsequently cleaved thereby. See Altman *et al.*, U.S. Pat. No. 5,168,053 and Yuan *et al.*, Science, 263:1269, (1994).

According to another preferred embodiment of the present invention the expression products of the first and second polynucleotides form a hetero-oligomer, e.g., a heterodimer. Such a hetero-oligomer can be, for example, an enzyme which displays cytotoxic activity. Thus, according this aspect of the present invention the first and second promoters are both stamen-specific promoters, as a result of which both polynucleotides are expressed and their polypeptide products assembled

in staminal tissue, such as for example anther tissue, so as to cause degeneration of anther tissue which ultimately leads to male sterility.

To generate exogenic allelism in plants any one of the expression cassettes described hereinabove must first be used to transform plant
5 tissue from which transgenic plants can be reconstituted.

As is further detailed above the term "transformation" describes a process by which an exogene such as an expression cassette enters and changes a recipient cell into a transformed, genetically modified or transgenic cell. Transformation may rely on any known method for the
10 insertion of foreign nucleic acid sequences into a eukaryotic host cell. The principle methods of effecting stable integration of exogenous DNA into plant genomic DNA include, Agrobacterium-mediated gene transfer, direct DNA uptake, including methods for direct uptake of DNA into protoplasts, electroporation of DNA, injection of DNA into plant cells or
15 tissues, biolistic bombardment of DNA coated particles, direct incubation of DNA with germinating pollen and the use of plant viruses as gene vectors. These methods are well described in the art and as such no further description is provided herein.

By applying any one of the methods for generating exogenic
allelism described hereinabove, resultant male sterile, female fertile
plants can be obtained. Such male sterile plants, when out-crossed with a
male fertile plant yield male fertile hybrid seeds in 100 % of the progeny
5 obtained.

Thus, two exogenous and distinct polynucleotide sequences which
exhibit allelic relationship, can be co-expressed within the same
transgenic plant and as a result of which induce male sterility.
Subsequent segregation of these sequences leads to recovery of fertility
10 in 100 % of the cases. Such segregation will be observed in 100 % of the
offsprings when such a plant is crossed with a compatible non-transgenic
plant. As such, exogenic allelism can be used to generate male sterility in
plants, which male sterility is 100 % reversible in subsequent
generations. This is of particular advantage since recovery of fertility is
15 of particular importance in hybrid crop production in which seeds or
fruits are collected. Such complete recovery of fertility can only be
achieved by complete segregation of the two polynucleotide sequences
and as such it can only be achieved for sequences which are disposed
within the genome of the plant in allelic relationship. As such, complete

segregation and as a result complete recovery of fertility are ensured by the methods and expression cassettes of the present invention.

It will be appreciated that the methods and constructs of the present invention can be used to generate reversible male sterility in both
5 fruit and non-fruit producing plants. Example of fruit producing plants include but are not limited to, wheat, barley and rice, while examples to non-fruit producing plants include, but are not limited to, lettuce, cauliflower and cabbage. Of particular importance are plants from which
10 fruits are harvested such as grains, since the resultant hybrid seeds must be male fertile such that plants grown from such seeds bear fruit which can be harvested. In this case 100 % recovery of fertility is of economic importance since all of the resultant progeny from the hybrid cross will produce fruit. It will be appreciated by one of skill in the art that the expression cassettes and methods described herein can also be applied to
15 *animalia*.

For example, exogenic allelism can be used to express equal-molar amounts of two distinct exogenes which may otherwise be subjected to differential positional effects. For example a cell culture derived from cell of an animal characterized by exogenic allelism can be

used to express two distinct exogenes in substantially equal-molar amounts since both exogenes are positioned in an identical position on two chromosomes of a chromosome pair. As a result these exogenes will be subjected to the same positional effect if any and as such will be expressed in substantially equal-molar amounts. It will be appreciated in this case that identical promoter sequences are employed upstream to the two exogenic sequences.

It will be appreciated that methods of animal cell transformation and generation of transgenic animals are well known in the art. Such methods are further described in the manual and text books listed in the preamble of the Examples section below.

Thus, the present invention provides methods and expression cassettes with which exogenic allelism can be generated in a non-human eukaryotic organism. Such allelism can be of great utility in cases where complete segregation of the allelic exogenes is desirable in subgenerations, such as the case of male sterile plants and their resultant male fertile hybrid progeny.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Cell Biology: A Laboratory Handbook" Volumes I-III Cellis,

J. E., ed. (1994); "Current Protocols in Immunology" Volumes I-III
Coligan J. E., ed. (1994); "Oligonucleotide Synthesis" Gait, M. J., ed.
(1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J.,
eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins
5 S. J., eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986);
"Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide
to Molecular Cloning" Perbal, B., (1984) and "Methods and
Enzymology" Vol. 1-317 Academic Press; all of which are incorporated
by reference as if fully set forth herein. Other general references are
10 provided throughout this document. The procedures therein are believed
to be well known in the art and are provided for the convenience of the
reader. All the information contained therein is incorporated herein by
reference.

EXAMPLE 1

15 Example 1 describes an expression cassette suitable for the
generation of allelism in a plant, which allelism is used to generate 100 %
reversible male sterility. Since hybrid seed production is of commercial
importance and since complete recovery of male fertility is crucial for
high grain crop yields a system was developed which allows for the

generation of male sterility but at the same time allows complete recovery of male fertility in progeny resultant from an outcross such as, for example, a hybrid outcross. As such this system can be used with high efficiency in hybrid seed production.

5 It will be appreciated that methods pertaining to plant self crossing and outcrossing utilized herein for the propagation of transformed plants and for subsequent hybrid seed production are commonly practiced in the art and as such no further description is necessary.

 Figure 1 outlines one method for generating reversible male
10 sterile plants according to the teachings of the present invention.

 Plant tissue is transformed with cassette I by, for example, an agrobacterium mediated gene transfer method and transgenic tissue is regenerated into identical transgenic plants which harbor a single copy of cassette I. Cassette I includes a Tapetum specific promoter (TapP) such
15 as ant32 or any other suitable stamen tissue specific promoter, followed by a LOX direct repeat. Following the LOX sequence is a T7 promoter region (T7P) which is recognized by the bacteriophage T7 polymerase which is not normally produced in plants. As such expression from the T7 promoter requires the presence of exogenous T7 polymerase gene or

gene product. The T7 promoter is operatively linked to a cell cytotoxic molecule coding sequence designated as the toxin. In the presence of a T7 polymerase the toxin is expressed from the T7 promoter. A toxin which can be used in accordance with the present invention encodes a protein or an RNA molecule capable of disrupting the production of functional pollen cells or the formation of pollen as further detailed hereinabove. The toxin gene is followed by a second Lox site (direct repeat) which is followed by a T7 polymerase gene which is not linked to an operable promoter.

The transgenic plant including this cassette is crossed with an isogenic plant which is homozygous for a recombinase gene (CRE) expressible under a strong constitutive promoter such as the 35 S promoter but which does not contain cassette I. The transgenic plant including cassette I is also selfed to obtain a homozygote (III).

The plant resultant from cross II is selfed to obtain a plant containing cassette IV. Cassette IV results from the excision of the toxin and the T7 promoter regions. This excision brings the T7 polymerase region in close proximity to the TapP promoter region (separated by the Lox site which is 34 bp in length) such that the T7 polymerase gene is

now under the transcriptional control of the TapP promoter. The plants harboring cassette IV will express the T7 polymerase in tapetal cells which should have no deleterious effects on pollen production. It will be appreciated in this case that since the direct repeat (Lox) site is immediately downstream of the TapP promoter it is important to either use direct repeats which do not contain an ATG codon therein or to eliminate any possible ATG start codon from the direct repeat sequence (using site specific mutation) such that transcription starts from the T7 polymerase ATG.

In the next step plant IV is crossed with plant III which does not express any of its exogenes, to yield a plant including a chromosome pair which is made up of a chromosome from plant III and a chromosome of plant IV. The resultant plant (V) is male sterile because the T7 polymerase produced in tapetal cells from chromosome A binds the T7 promoter region on chromosome A and drives the expression of the toxin which leads to the degeneration of the tapetal cells and to male sterility.

This plant serves as the maternal plant for subsequent hybrid crosses (VII) in which the pollen of a compatible non-transgenic male fertile plant (VI) is used to fertilize plant V to yield (from plant V) fertile

offsprings. This is achieved because of the 100 % segregation of chromosomes A and A. The seeds collected from plant V are hybrid seeds which include either cassette VIIa or cassette VIIb.

It will be appreciated that the specific arrangement of the genes,
 5 recombination sites and promoters in cassette I can be altered and yet produce very similar results. For example the TapP promoter can be linked to the T7 polymerase and both flanked by the Lox sites. In this case the T7 promoter can be separated from the toxin gene by the TapP promoter and the T7 polymerase. Following excision the toxin is brought
 10 under the control of the T7 promoter.

EXAMPLE 2

An alternative method for generating allelism is shown in Figure
 2. Again this method is shown with specific reference to induced male
 15 sterility, it is to be understood however that his method can be applied to generate allelism for any purpose.

Figure 2 outlines another method for generating reversible male sterile plants according to the teachings of the present invention.

Plant tissue is transformed with cassette I and transgenic tissue is regenerated into identical transgenic plants. These transgenic plants are male sterile because they express both the T7 polymerase and as a result the toxin expressed from the T7 promoter

5 According to this method cassette I includes a T7 polymerase operatively linked to the TapP promoter such as, for example, ant32, both flanked by the FRT recombinase sites. Cassette I further includes a toxin gene operatively linked to a T7 promoter, both flanked by the Lox sites.

10 Identical plants harboring cassette I are pollinated by a Cre recombinase plant (II) (homozygous for Cre) and an FLP recombinase plant (III) (homozygous for FLP). The resultant plants (IV and V) which contain either one of the gene and linked promoter are male fertile. As such these plants are selfed in order to lose the recombinase gene and to establish homozygotes of their recassettes (VI and VII). Plants VI and 15 VII are crossed and the generated male sterile plant includes the two exogenes (T7 polymerase and the Toxin) in an allelic relationship. From this point on production of hybrid male fertile plants follows that described in Example 1.

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---